ourdiu.com | Shared by Md. Shefat Al Mahmud

Daffodil International University

Department of Computer Science and Engineering
Faculty of Science & Information Technology
Semester Final Examination, Spring 2023
Course Code: STA 221, Course Title: Statistics and Probability
Level: 2 Term: 2 Batch: All

Time: 02:00 Hrs **Marks: 40**

Answer ALL Questions

[The figures in the right margin indicate the full marks and corresponding course outcomes. All portions of each question must be answered sequentially.]

1.	*a)*	If you want to calculate the test scores on a final exam to better understand whether most students score close to the average or if there is a wide spread in test scores. **Find** the suitable measure of dispersion for the analysis and **explain**.	[1]	CO1
	b)	**Show** the difference mutually exclusive and independent events? Illustrate with a hypothetical example.	[2]	
	c)	Meditation can produce a deep state of relaxation and a tranquil mind. During meditation, you focus your attention and eliminate the stream of jumbled thoughts that may be crowding your mind and causing stress. **Choose** the appropriate correlation is existing in this example. Explain	[2]	
2.		Software firm collected data for a sample of 9 programmer's firm's management wanted to predict salary using years of experience. The data are given below:		CO2

Salary(K)	24	43	47	51	35	38	42	40	49
No. of experience (years)	4	7	6	7	5	5	6	5	6

	i)	**Construct** a scatter diagram.	[2]
	ii)	**Develop** coefficient of determination using the above table.	[4]
	iii)	Test the hypothesis that there is a significant relationship between salary and **No. of experience (years)** used at the 0.05 significance level, **interpret** your findings. [Tabulated Value: 1.86]	[5]
3.	*a)*	By using the data 2		CO3
		i) **Build** a regression model and interpret the results	[5]	
		ii) **Build** salary if number of experiences is 10 years	[1]	
	b)	If two dice are thrown **solve** the probabilities that the both would be odd, same number?	[3]	
4.	*a)*	The number of hits on a certain website follows poison distribution with a mean rate of 4 per minutes. **Discover**		CO4
		i) the probability that at least 2 hits are received in 1 minutes	[2.5]	
		ii) the probability that between 2 to 3 hits are received in 2 minutes	[2.5]	
	b)	A package filling process at a cement fills a bag of cement to an average weight of 12 pounds, the standard deviation is 3 pounds. A sample of 15 bags have been taken and the mean was found to be 100 pounds. Assume that the weights of the bag follow normal distribution, and **test** the above process at 5% level of significance. [Z cal.=1.96]	[5]	
	c)	It is observed that 50% of mails are spam. There is a software that filters spam mail before reaching the inbox. Its accuracy for detecting a spam mail is 99% and chances of tagging a non-spam mail as spam mail is 5%. If a certain mail is tagged as spam **discover** the probability that it is not a spam mail.	[5]	

ourdiu.com | Shared by Md. Shefat Al Mahmud

Formulas

Measure of Dispersion

Range= X_{max} - X_{min}

Mean Deviation, M.D $=\frac{\sum_{i=1}^{n}|x_i-\bar{x}|}{n}$

Population variance

$\sigma^2=\frac{\sum_{i=1}^{N}(X_i-\mu)^2}{N}$

Sample variance

$s^2=\frac{\sum_{i=1}^{N}(X_i-\bar{X})^2}{n-1}$

Coefficient of variation for population, C.V$=\frac{\sigma}{\mu}\times 100$

Coefficient of variation for sample, C.V$=\frac{s}{\bar{x}}\times 100$

Basic Concepts of Probability

General Rule of Addition

P(A or B)= P(A U B)= P(A) + P(B) – P(A∩B)

Special Rule of Addition

P (A or B) = P (A U B) = P(A) + P(B)

Conditional Probability

P(B|A) = P(A and B) / P(A)= $\frac{P(A\cap B)}{P(A)}$,

Bayes theorem:

$$P[B_i|A]=\frac{P[B_i]P[A|B_i]}{\sum_{j=1}^{n}P[B_j]P[A|B_j]};\ i=1,2,\ldots\ldots,n$$

Hypothesis testing

Z test statistics $Z=\frac{\bar{x}-\mu}{\frac{\sigma}{\sqrt{n}}}$

t test statistics $t=\frac{\bar{X}-\mu}{s/\sqrt{n}}$

Testing the significance of the correlation coefficient

$$t=\frac{r}{\sqrt{\frac{1-r^2}{n-2}}}$$

Correlation and Regression analysis

Regression Coefficient

$$\hat{\beta}_1=\frac{\sum_{i=1}^{n}(x_i-\bar{x})(y_i-\bar{y})}{\sum_{i=1}^{n}(x_i-\bar{x})^2}$$

$$\hat{\beta}_0=\bar{y}-\hat{\beta}_1\bar{x}$$

Correlation Coefficient

$$r=\frac{N\Sigma xy-(\Sigma x)(\Sigma y)}{\sqrt{[N\Sigma x^2-(\Sigma x)^2][N\Sigma y^2-(\Sigma y)^2]}}$$

$$r=\frac{\sum_{i=1}^{n}(X_i-\bar{X})(Y_i-\bar{Y})}{\sqrt{\sum_{i=1}^{n}(X_i-\bar{X})^2}\sqrt{\sum_{i=1}^{n}(Y_i-\bar{Y})^2}}$$

Binomial Distribution

$$f(x;n,p)=\begin{cases}\binom{n}{x}p^x q^{n-x} & \text{for } x=0,1,2,\ldots,n\\ 0; & \text{otherwise}\end{cases}$$

Poison Distribution

$$f(x;\lambda)=\begin{cases}\frac{e^{-\lambda}\lambda^x}{x!} & \text{for } x=0,1,2,\ldots,\infty.\\ 0; & \text{otherwise}\end{cases}$$

Normal Distribution

$$f(x,\mu,\sigma^2)=\frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{(x-\mu)^2}{2\sigma^2}};\qquad -\infty<x<\infty$$

Shape of the distribution

Coefficient of Sknewness,

Sk$=\frac{3\times(Mean-Median)}{Standard\ deviation}$

Kurtosis $\beta_2=\frac{\mu_4}{\mu_2^2}$